

TECHNICAL SHEET IDYLLA™ BRAF MUTATION ASSAY

The Idylla™ BRAF Mutation Assay, performed on the Biocartis Idylla™ System, is a molecular assay for the qualitative detection of V600E/E2/D and V600K/R/M mutations in codon 600 of the BRAF gene.

The Idylla™ BRAF Mutation Assay, from sample-to-result, starts with formalin-fixed paraffin-embedded (FFPE) tissue to liberate DNA for subsequent real-time PCR amplification and detection.

FEATURES

BRAF mutation detection		
	BRAF V600E BRAF V600E2 BRAF V600D	(c.1799T>A) (c.1799_1800delinsAA) (c.1799_1800delinsAT; c.1799_1800delinsAC)
Codon 600	BRAF V600K BRAF V600R BRAF V600M	(c.1798_1799delinsAA) (c.1798_1799delinsAG) (c.1798G>A)
	BRAF Wild Type	(c.1799T)

RNaseP (acting as Sample Processing Control)

RNaseP (acting as Sample Processing Control)		
Specimen requirements		
Sample Type	FFPE tissue sections (5 to 10 μm)	
Neoplastic cells	≥50%, if less macrodissection is required	
Tissue area	50-600 mm² (5 μm) 25-300 mm² (10 μm)	
Total turnaround time		
Time	90 minutes	
Result Reporting		
Report	Qualitative genotype call	



