

NOTHING
IS SIMPLE
IN ONCOLOGY.

NOTHING BUT THIS.




Idylla[™] A revolutionary, fully automated system that makes molecular testing convenient and fast. Suitable for any lab. biocartis.com/us



WE AIM TO PROVIDE RAPID ACCESS TO HIGH-LEVEL MOLECULAR TESTING ANYTIME, ANYWHERE.

THE IDYLLA™ SOLUTION - MOLECULAR TESTING SUITABLE FOR VIRTUALLY ANY LAB.



^{*}For Idylla™ IVD Tests.

IDYLLA™ FITS ANY SIZE LAB



IDYLLA™ IS A COST-EFFECTIVE IN-HOUSE SOLUTION EVEN FOR SMALL SAMPLE VOLUMES

- Easy to implement and maintain
- No molecular infrastructure needed
- No batching needed




IDYLLA™ PROVIDES AN ULTRA-RAPID ACTIONABLE SOLUTION COMPLEMENTING COMPREHENSIVE NGS PROGRAMS

- Short turnaround time and minimal hands-on time
- Minimal sample input requirements
- On-demand testing
- Seamless integration into virtually any laboratory workflow




THE REVOLUTIONARY IDYLLA™ WORKFLOW

STANDARD SOLID BIOPSY WORKFLOW



LIQUID BIOPSY WORKFLOW




5

IDYLLA™ COMPARED TO OTHER TECHNOLOGIES

MINIMAL HANDS-ON AND ASSAY TURNAROUND TIMES



REDUCED NUMBER OF INSTRUMENTS AND CONSUMABLES NEEDED



Based on workflow exercise in real-life laboratory setting. Reference technologies used: Illumina $MiSeq^*$ and $Qiagen\ Therascreen^*$.

THE IDYLLATM ASSAYS. MINIMAL SAMPLE REQUIREMENTS AND VERSATILE SAMPLE TYPES.



SOLID BIOPSY SPECIMEN REQUIREMENTS

IVD TESTS

MSI (510(k))	 1 x 4 µm FFPE tissue section 1 x 5 µm FFPE tissue section 1 x 10 µm FFPE tissue section Neoplastic cells ≥ 33% - if less, macrodissection is required 	\rightarrow 62.5-750 mm ² \rightarrow 50-600 mm ² \rightarrow 25-300 mm ²
MSI-CDx (PMA) RUO ASSAYS	 1 x 4 μm FFPE tissue section 1 x 5 μm FFPE tissue section 1 x 10 μm FFPE tissue section Neoplastic cells ≥ 33% - if less, macrodissection is required 	\rightarrow 62.5-750 mm ² \rightarrow 50-600 mm ² \rightarrow 25-300 mm ²
EGFR	 1 x 5 µm FFPE tissue section Neoplastic cells ≥ 10% - if less, macrodissection is required 	→ ≥1 mm²
GeneFusion	 1 x 5 µm FFPE tissue section 3 x 5 µm FFPE tissue sections Neoplastic cells ≥ 10% - if less, macrodissection is required 	\rightarrow 20-100 mm ² \rightarrow < 20 mm ²
BRAF	 1 x 5 µm FFPE tissue section 1 x 10 µm FFPE tissue section Neoplastic cells ≥ 50% - if less, macrodissection is required 	\rightarrow 50-600 mm ² \rightarrow 25-300 mm ²
KRAS, NRAS-BRAF-EGFR S492R	 1 x 5 µm FFPE tissue section 1 x 10 µm FFPE tissue section Neoplastic cells ≥ 10% - if less, macrodissection is required 	→ 50-600 mm² → 25-300 mm²
POLE-POLD1	 1 x 5 µm FFPE tissue section 1 x 10 µm FFPE tissue section Neoplastic cells ≥ 10% - if less, macrodissection is required 	→ 50-600 mm² → 25-300 mm²
IDH1-2	 50 μl extracted DNA (concentration > 10 ng/μl) 10 μl blood or bone marrow Maximum 3 FFPE tissue sections Neoplastic cells > 10% - if less, macrodissection is required 	
PIK3CA-AKT1	 1 x 5 μm FFPE tissue section 1 x 10 μm FFPE tissue section Neoplastic cells ≥ 20% - if less, macrodissection is required 	→ 50-600 mm ² → 25-300 mm ²



LIQUID BIOPSY SPECIMEN REQUIREMENTS

RUO ASSAYS

• 1 or 2 ml plasma (depending on Assay) and two validated tube types:



K²EDTA tubes

process within 4 hours



Streck Cell-Free DNA BCT® Tubes

process within 3 days

• Used in the following Idylla™ Assays: Idylla™ ctKRAS, ctEGFR and ctNRAS-BRAF-EGFR S492R Mutation Assays (RUOs)



CONTACT



Biocartis US Inc. 2 Pierce Place, Suite 1510 Itasca, IL 60143 - US +1 (844) 443-9552 REQUEST A DEMO



Follow us on **f X in** biocartis.com/us customerserviceUS@biocartis.com

